

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	1498	(544/281,319).CCLS.	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	OFF	2004/09/16 11:15
L2	1871	((544/3) or (514/222.2,259.1,269)). CCLS.	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	OFF	2004/09/16 11:15
L3	3000	1 or 2	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	ON	2004/09/16 11:16





Hye<sup>2</sup>

Hye<sup>3</sup>

Hye<sup>4</sup>

Hye<sup>5</sup>




19e<sup>2</sup>

27e<sup>3</sup>

28e<sup>4</sup>

29e<sup>5</sup>



ain nodes :

7 8 9 10 11 12 19 24 25 27 28 29

ng nodes :

1 2 3 4 5 6 13 14 15 16 17 18

ain bonds :

1-25 3-9 4-7 5-8 9-10 9-11 11-12 12-24

ng bonds :

1-2 1-6 2-3 3-4 4-5 5-6 13-14 13-18 14-15 15-16 16-17 17-18

act/norm bonds :

1-2 1-6 1-25 2-3 3-4 4-5 4-7 5-6 5-8 9-10 9-11 11-12 12-24

act bonds :

3-9

rmalized bonds :

13-14 13-18 14-15 15-16 16-17 17-18

olated ring systems :

containing 1 :

:[\*1],[\*2]

:[\*1],[\*2],[\*3],[\*4],[\*5]

atch level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS  
11:CLASS 12:CLASS 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom 24:CLASS  
25:Atom 27:Atom 28:Atom 29:Atom

neric attributes :

19:  
Saturation : Unsaturated  
Number of Carbon Atoms : less than 7  
Number of Hetero Atoms : less than 2  
Type of Ring System : Monocyclic  
27:  
Saturation : Unsaturated

Number of Carbon Atoms : less than 7  
Number of Hetero Atoms : less than 2  
Type of Ring System : Monocyclic  
28:  
Saturation : Unsaturated  
Number of Carbon Atoms : less than 7  
Number of Hetero Atoms : less than 2  
Type of Ring System : Monocyclic  
29:  
Saturation : Unsaturated  
Number of Hetero Atoms : 2 or more  
Type of Ring System : Polycyclic

ement Count :

Node 19: Limited

C,C5  
N,N1  
O,O0  
S,S0

Node 27: Limited



C,C4  
N,N1  
O,O0  
S,S0

Node 28: Limited

C,C4  
O,O1  
N,N0  
S,S0

Node 29: Limited

N,N3  
C,C6  
O,O0  
S,S0



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Box  
Search PubMed for hiv integrase inhibitors treatment infection Go Clear

Limits Preview/Index History Clipboard Details

Display Summary Show: 20 Sort Send to Text

About Entrez

Items 1 - 13 of 13

One page

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

- ☐ 1: [Hazuda DJ, Young SD, Guare JP, Anthony NJ, Gomez RP, Wai JS, Vacca JP, Handt L, Motzel SL, Klein HJ, Dornadula G, Danovich RM, Witmer MV, Wilson KA, Tussey L, Schleif WA, Gabryelski LS, Jin L, Miller MD, Casimiro DR, Emini EA, Shiver JW.](#) Related Articles, Link

Integrase inhibitors and cellular immunity suppress retroviral replication in rhesus macaques.  
Science. 2004 Jul 23;305(5683):528-32. Epub 2004 Jul 08.  
PMID: 15247437 [PubMed - indexed for MEDLINE]

- ☐ 2: [de Soultrait VR, Desjobert C, Tarrago-Litvak L.](#) Related Articles, Link

Peptides as new inhibitors of HIV-1 reverse transcriptase and integrase.  
Curr Med Chem. 2003 Sep;10(18):1765-78. Review.  
PMID: 12871103 [PubMed - indexed for MEDLINE]

- ☐ 3: [Billich A.](#) Related Articles, Link

S-1360 Shionogi-GlaxoSmithKline.  
Curr Opin Investig Drugs. 2003 Feb;4(2):206-9. Review.  
PMID: 12669383 [PubMed - indexed for MEDLINE]

- ☐ 4: [Gulick RM.](#) Related Articles, Link

New antiretroviral drugs.  
Clin Microbiol Infect. 2003 Mar;9(3):186-93. Review.  
PMID: 12667250 [PubMed - indexed for MEDLINE]

- ☐ 5: [Garvey EP.](#) Related Articles, Link

Unexploited viral and host targets for the treatment of human immunodeficiency virus type 1 infection.  
Curr Drug Targets Infect Disord. 2001 Aug;1(2):107-23. Review.  
PMID: 12455408 [PubMed - indexed for MEDLINE]

- ☐ 6: [Chaisson RE, Gebo K, Flexner C, Gallant JE, Lucas GM, Bartlett JG.](#) Related Articles, Link


Selected topics from the 9th Conference on Retroviruses and Opportunistic Infections.  
HIV Clin Trials. 2002 Jul-Aug;3(4):333-49. No abstract available.  
PMID: 12187508 [PubMed - indexed for MEDLINE]


- ☐ 7: [Debyser Z, Cherepanov P, Van Maele B, De Clercq E, Witvrouw M.](#) Related Articles, Link

In search of authentic inhibitors of HIV-1 integration.  
Antivir Chem Chemother. 2002 Jan;13(1):1-15. Review.  
PMID: 12180645 [PubMed - indexed for MEDLINE]


- ☐ 8: [Tarrago-Litvak L, Andreola ML, Fournier M, Nevinsky GA, Parissi V, de Soultrait VR, Litvak S.](#) Related Articles, Link

Inhibitors of HIV-1 reverse transcriptase and integrase: classical and

 emerging therapeutical approaches.  
Curr Pharm Des. 2002;8(8):595-614. Review.  
PMID: 11945161 [PubMed - indexed for MEDLINE]


 **9:** [Raulin J.](#)


[Related Articles](#), [Link](#)

 Human immunodeficiency virus and host cell lipids. Interesting pathways in research for a new HIV therapy.  
Prog Lipid Res. 2002 Jan;41(1):27-65. Review.  
PMID: 11694268 [PubMed - indexed for MEDLINE]


 **10:** [Jarmy G, Heinkelein M, Weissbrich B, Jassoy C, Rethwilm A.](#)


[Related Articles](#), [Link](#)

 Phenotypic analysis of the sensitivity of HIV-1 to inhibitors of the reverse transcriptase, protease, and integrase using a self-inactivating virus vector system.  
J Med Virol. 2001 Jul;64(3):223-31.  
PMID: 11424108 [PubMed - indexed for MEDLINE]


 **11:** [Whitson S.](#)


[Related Articles](#), [Link](#)

 Drug watch.  
Posit Aware. 1999 Mar-Apr;10(2):14-5.  
PMID: 11366839 [PubMed - indexed for MEDLINE]


 **12:** [Pani A, Marongiu ME.](#)

[Related Articles](#), [Link](#)

 Anti-HIV-1 integrase drugs: how far from the shelf?  
Curr Pharm Des. 2000 Mar;6(5):569-84. Review.  
PMID: 10788597 [PubMed - indexed for MEDLINE]

 **13:** [McDougall B, King PJ, Wu BW, Hostomsky Z, Reinecke MG, Robinson WE Jr.](#)

[Related Articles](#), [Link](#)

 Dicafeoylquinic and dicafeoyltartaric acids are selective inhibitors of human immunodeficiency virus type 1 integrase.  
Antimicrob Agents Chemother. 1998 Jan;42(1):140-6.  
PMID: 9449274 [PubMed - indexed for MEDLINE]

[Display](#)

[Summary](#)

Show: [20](#)

[Sort](#)

[Send to](#)

[Text](#)


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Sep 10 2004 06:30:44



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Box

Search PubMed for

Limits Preview/Index History Clipboard Details

Abstract  20   Text

About Entrez

Text Version

☐ 1: Prog Lipid Res. 2002 Jan;41(1):27-65.

Related Articles, Lir

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

ELSEVIER SCIENCE  
FULL-TEXT ARTICLE

## Human immunodeficiency virus and host cell lipids. Interesting pathways in research for a new HIV therapy.

**Raulin J.**

Universite Denis Diderot (Paris 7), 2 place Jussieu, 75251 05, Paris, France.  
raulinj@hotmail.com

It has been reported in the literature that biological membranes arising from HIV-induced cell fusion, as well as syncytium formation between infected and non-infected cells and those involved in transduction, viral DNA nuclear import and virion budding from the host cell, are all made of proteins, a phospholipid (P) bilayer and cholesterol (C). However, the P/C molar ratio is higher in the retroviral envelope than in the plasma membrane where they originate, and higher than in the nuclear envelope. Mechanisms are described which elucidate this puzzling fact, as well as cholesterol-dependent leakage and pore formation during cell fusion. Fatty acylation of viral and host cell proteins is required to direct them to membranes. Detergent-insoluble microdomains enriched in cholesterol and sphingolipids, termed either DIGs (detergent-insoluble glycolipid-enriched complexes), DRMs (detergent resistant membranes), TIFFs (Triton-insoluble floating fractions) or GEMs (glycolipid-enriched membranes), function as platforms for attachment of proteins in the process of signal transduction. HIV-SUgp120 (HIV-surface glycoprotein), T-cell receptor (TCR)-CD4+ and co-receptors promote aggregation of these lipid "rafts" which concentrate the Src family tyrosine kinases SFKs (PTK, Lyn, Fyn, Lck), GPI (glycosyl phosphatidylinositol)-anchored proteins, and phosphatidylinositol kinases PI(3)K and PI(4)K, inducing cell signalling. HIV-SUgp120 transduces the activation signal and provokes the formation of polyunsaturated fatty acid (PUFA) metabolites, i.e. the prostaglandin PGE2 suppressor of immune function and inhibitor of cytotoxic T-lymphocyte (CTL) proliferation, while PGB2 activates SFKs and increases mRNA expression, as well as NFkappaB (nuclear transcription factor) translocation to nucleus. HIV nuclear import, DNA integration, chromatin template capacity may be mediated by the lipid environment. The lipid-enriched microdomains from which HIV-1 buds, may explain the high level of cholesterol and sphingolipids in the viral envelope, since host cell rafts become a viral coat. HIV-1 infection induces alteration of cellular lipids (1) shift in phospholipid synthesis to neutral lipids associated with the viral

load, polyunsaturated fatty acid (PUFA) peroxidation, and n-3 deficiency with deregulation of cytokines and PPAR-gamma (peroxisome proliferator-activated receptor-gamma), and (2) alloimmune phospholipid antibody production in which antibodies to cardiolipin and to phosphatidylserine are most prevalent, due to the destruction of mitochondrial membranes and progression of lymphocyte apoptosis. The current highly active anti-retroviral therapy, including both viral reverse transcriptase (RT) inhibitors (NRTIs and NNRTIs, nucleoside and non-nucleoside RT inhibitors) and protease inhibitors (PIs), induces side-effects in the long term. Lipodystrophy (LD), consists of peripheral lipoatrophy associated with central fat accumulation (called "crixibelly" and "buffalo hump"), insulin resistance, elevation of very low density lipoproteins, decrease in high density lipoproteins and inhibition of adipocyte differentiation. LD syndrome appears to be induced by PIs that inhibit GLUT4, glucose transporter isoform, and by NRTIs which provoke mitochondrial failure. New therapeutic strategies assessed: (1) inhibition of viral integrase and/or HIV entry into cells through natural products or their derivatives, (2) inhibition of HIV-1 entry into macrophages pretreated with Gram-negative bacterial lipopolysaccharide, (3) vaccination with multi-lipopeptides, i.e. sequences of HIV-1 peptides with CD4+ T-cell and B-cell epitopes, modified by adding a lipid tail to one end, which produce HIV-specific CTL and multispecific immune responses in most of the vaccinated subjects and (4) stimulation of antiviral drug activity with lipid-prodrugs targeting viral RT, polymerase, integrase, or aspartyl-protease.

Publication Types:

- Review
- Review, Academic

PMID: 11694268 [PubMed - indexed for MEDLINE]

Display

Abstract



Show:

20



Sort



Send to

Text



[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Sep 10 2004 06:30:44